



ARTICLE

Comparison of Functional Compounds and Micronutrients of Chicken Breast Meat by Breeds

Mahabbat Ali¹, Seong-Yun Lee¹, Ji-Young Park¹, Samooel Jung², Cheorun Jo³, and Ki-Chang Nam^{1,*}

¹Department of Animal Science and Technology, Suncheon National University, Suncheon 57922, Korea

²Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Korea

³Center for Food and Bioconvergence, and Research Institute for Agriculture and Life Science, Seoul National University, Seoul 08826, Korea

OPEN ACCESS

Received June 12, 2019

Revised June 28, 2019

Accepted July 18, 2019

*Corresponding author : Ki-Chang Nam
Department of Animal Science & Technology,
Suncheon National University, Suncheon
57922, Korea
Tel: +82-61-750-3231
Fax: +82-61-750-3231
E-mail: kichang@scnu.kr

*ORCID
Mahabbat Ali
<https://orcid.org/0000-0002-5332-7758>
Seong-Yun Lee
<https://orcid.org/0000-0003-1361-1962>
Ji-Young Park
<https://orcid.org/0000-0001-9680-8685>
Samooel Jung
<https://orcid.org/0000-0002-8116-188X>
Cheorun Jo
<https://orcid.org/0000-0003-2109-3798>
Ki-Chang Nam
<https://orcid.org/0000-0002-2432-3045>

Abstract The concentrations of functional compounds and micronutrients of chicken breast from native chickens were compared with those from broiler. Totally 200 male chicks from a commercial native chicken (HH) and three newly bred native chicken strains (2A, 2C, and 2D) were reared for about 2 kg of final live weight up to 12 wk. After slaughtered, antioxidant dipeptides, reducing sugar, free amino acids, vitamins, and minerals of the breast muscles were analyzed with those from broilers with similar live weight. Mostly native chicken strains had higher contents of carnosine, anserine, and reducing sugar than the broiler. Especially HH implied the highest values of carnosine and anserine, and 2C did the highest of reducing sugar ($p < 0.05$). Vitamin A contents between native chickens and broiler were not significantly different ($p > 0.05$). The contents of α -tocopherol were significantly higher in 2C than those of HH or broiler ($p < 0.05$). Native chicken strains contained lower cholesterol levels than the broiler. Broiler had higher contents of P, Mg, and Na than native chickens ($p < 0.05$), but it had lower content of Cu than HH or 2A. The savory free amino acids including glutamic acid was highest in 2A than the other native chickens and broiler ($p < 0.05$). This study confirms that certain new strains of native chickens be a good source in terms of functional compounds and micronutrients which can be attractive health promoting nutritional quality factors.

Keywords antioxidant dipeptides, reducing sugar, functional compounds, micronutrients, chicken strain

Introduction

Chicken meat is widely consumed in Southeast Asia considering as traditional healthy food (Nagai et al., 1996). Over the past few decades the per capita of poultry meat has been increased by approximately 3 kg based on report of the Korean Ministry of Agriculture and Forestry. Despite of increasing circumstances, the Korean native

chickens have not been produced sufficient amount because of the limited productivity with desirable characters of meat.

Some bioactive endogenous compounds such as carnosine (β -alanyl histidine), anserine (1-methyl carnosine) in the skeletal muscle of vertebrate animals was considered as strategy for poultry production (Schmid, 2009). Mammalian skeletal muscles rich with carnosine act as a pH buffer in muscles, antiglycation, antiaging, antioxidation and neurotransmitter functions (Schmid, 2009). The color and flavor of meat depends on oxidative mechanisms (Ouali et al., 2006). Carnosine combination with dietary vitamin E supplementation improves the lipid stability in meat (Morrissey et al., 1998). Carnosine also suppresses some diseases such as diabetes, cataracts, ischemia, and Alzheimer's diseases (Hipkiss, 2009) and attenuates acidosis as well as loss of force during exercise in human body (Baguet et al., 2010). Carnosine is an activator of several enzymes including calpain II, myofibrillar ATPase and phosphorylase (Iwaniak and Dziuba, 2009). Anserine (1-methylated carnosine) derivate from carnosine perform similar functions as like carnosine (Schmid, 2009).

Reducing sugars are important addition in meat in term of flavor react with amino acids by Maillard reaction during cooking (Mottram and Maarse, 1991). Maillard reaction occurs between an amino compounds (amine, amino acid, peptide, or protein) and a carbonyl compounds (usually a reducing sugar), the thermal degradation of thiamin, the oxidation of lipids, and the interactions between these pathways. The reaction of sugar and cysteine can lead to characteristic meat flavour specially for chicken (Varavinit et al., 2000). This was further confirmed by a research where the quantities of carbohydrates and amino acids, in particular ribose and cysteine, are reduced during heating. The main carbohydrates with flavour-forming potential include ribose, ribose-5-phosphate, glucose and glucose-6-phosphate (Meinert et al., 2009). The addition of small quantities of ribose in chicken meat has been shown to enhance the quantities of key odor compound as well as meaty and roasted flavor (Aliani and Farmer, 2005).

Containing all essential amino acids, fatty acids, vitamins, mainly B complex vitamins and especially B₁₂; minerals, mainly iron and zinc and other bioactive compounds meat and meat products are very familiar as high nutritious foods (WHO, 2003). Free amino acids, nucleic acids, peptides, and minerals are the compounds involved in meat flavor (Yamaguchi, 1991). Dietary cholesterol is strongly associated with coronary heart disease and arteriosclerosis that causes human mortality (Simopoulos, 2004). Recently many attempts has been taken to decrease the cholesterol contents of chicken meat adding dietary supplements like omega-3, garlic, and copper (Chowdhury and Smith, 2002). Chicken meat is a great source of many functional compounds and micronutrients i.e. iron, vitamin B₁₂ as well as zinc, B complex, selenium and phosphorus (Pereira and Vicente, 2013). Chicken meat contains of many minerals (Zn, Fe, and Cu) are essential for life and also may attribute for consumer satisfaction (Ribeiro et al., 2019).

Recently attempts has been taken by Korean government to develop new breed of chicken. This project mostly focused on developing the meat producing chicken breed with high quality traits by using Korean native chickens in order to meet the consumers demands (Park et al., 2012). Three new native chicken strains (i.e., 2A, 2C, and 2D) were anticipated as candidates for selection considering their growth rate and eating qualities. The aim of this study was to investigate the concentration of health promoting compounds such as antioxidant dipeptides, reducing sugar, and micronutrients contents of three new native chicken strains compared with commercial Korean native chicken and broiler to pick up the data which will be helpful for future development of breeding selection strategies for poultry meat production.

Materials and Methods

Birds

Totally 200 male chicks from a commercial native chicken (HH) and three newly bred native chicken strains (2A, 2C, and

2D) were reared up to 12 wk at an experimental farm of Harim corporation (Gimje, Korea) under similar conditions and diets. The new native chicken strains were bred by mating combinations from a paternal line (2) and 3 different maternal lines (A, C, and D). Broilers were also reared up to reach at similar slaughter weight with those HH and new native chicken strains. HH and broiler were considered as controls. Native chicken, three newly bred native chicken strains were slaughtered with those broiler with similar live weight, then carcass were vacuum-packed after chilling at 4°C for 24 h, and stored in a freezer at -20°C until analysis. The breast fillets were used for analysis.

Sample preparation

The frozen carcasses were thawed in a refrigerator at 4°C for 24 h before analysis in the laboratory. After thawing the right and left muscles were dissected and breast muscles also dissected in following pattern. After separation of samples then should be minced using food mixer. Then from minced meat samples, target supernatant were made in different terms and conditions to perform the respective analysis.

Reagents and materials

Standards (carnosine, and anserine) were obtained from Sigma-Aldrich (St. Louis, MO). The external and internal standard (IS) used in cholesterol analysis were cholesterol (Tokyo chemical industry, Tokyo, Japan) and 5 α -cholestane (Sigma-Aldrich Co., St. Louis, MO, USA). Standard products used for retinol and tocopherol analysis were purchased from Sigma-Aldrich Co. (St. Louis, MO, USA). Other reagents and solvents were used HPLC grade. Vitamin B₁₂ standards (cyanocobalamin, purity 96.0%–102.0%) were purchased from Sigma-Aldrich (St. Louis, Mo., USA) and sodium acetate trihydrate and acetic acid were purchased from Wako (Osaka, Japan).

Laboratory analysis

The contents of carnosine and anserine were determined using the slightly modification method ascribed by (Mora et al., 2007) where 2.5 g of minced meat samples from each chicken were homogenized with 7.5 mL of 0.01N HCl at 1,130 \times g for 30 s (T25 basic, IKA GmbH & Co. KG, Germany). After homogenization then the samples were centrifuged at 10,000 \times g for 30 min at 4°C (1580 R, GYROZEN Co., Ltd). 0.5 mL of centrifugated supernatant was taken in 2 mL tube and mixed with 1.5 mL of acetonitrile and performed centrifuge again for 10 min at 10,000 \times g at 4°C (1580 R, GYROZEN Co., Ltd). Filtrated through 0.2 μ m membrane filter the supernatant was injected into HPLC (1580R, GYROZEN Co., Ltd) column.

The reducing sugar contents of each extract were measured by dinitrosalicylic acid method as described by the (The Korean Society of Food Science and Nutrition, 2000). Beta-carotene content was determined by using HPLC (Agilent, Santa Clara, CA, USA) following the method of (Thomas et al., 2001) after extracting with alkali saponification method.

Tocopherols and tocotrienols were determined by mobile phase HPLC after saponification followed by (Food and Drug, 2016). 2 mL of the extract was taken accurately, the solvent was volatilized with nitrogen, and redissolved with 1 mL of n-hexane. The redissolved extract was filtered through a 0.5 μ m membrane filter (Advantec, Tokyo, Japan) and analyzed by HPLC (LC-20AD, Shimadzu, Kyoto, Japan). The column was detected using ExT 1 λ =285 nm and Em λ =325 nm using a Li Chrospher Diol 100 (240 \times 4 mm, 5 μ m, Merck, Darmstadt, Germany) using a fluorescence detector (Shimadzu).

Vitamin B₁₂ content was measured by the method of (Mun et al., 2017) equipped with HPLC-DAD analysis through immunoaffinity clean-up techniques where vitamin B₁₂ standard solution is prepared by dissolving 10 mg of cyanocobalamin standard product in 10 mL of water and making stock solution of 1 mg/mL. Then stored it in cold dark place at -18°C, diluted to

the range of 0.025–10 µg/mL. The column was incubated with the C18 ACE 3 AQ (3 mm×150 mm, ACE, Scotland, UK) using the HPLC system Agilent 1260 infinity (Agilent, Santa Clara, CA, USA) Gradient conditions. At this time, the flow rate of the mobile phase was 0.25 mL/min, the column oven temperature was 35°C, and the sample injection amount was 100 µL. The separated cyanocobalamin was detected at 361 nm, and the absorbance at 200–600 nm was collected to confirm the peak purity.

Minerals of each chicken breast fillet were analyzed by inductively coupled-plasma atomic emission spectrophotometry with the standard absorbance solutions. The cholesterol contents of minced samples were analyzed by GC (HP 5890 Series II, Hewlett Packard Co., Palo Alto, CA, USA). Analytical column was HP-5 (30 m×0.320 mm, 0.25 µm, Agilent, J & W Scientific, Santa Clara, CA, USA).

The soluble amino acids composition was determined by using a slightly modification of the method which is described by (Huge et al., 2002) where the visible external fat was removed and meat sample three grams from each treatment was mixed with 27 mL of 2% TCA solution. The mixture was homogenized at 13,500 rpm/min for 1 min. The homogenate was then centrifuged at 17,000 g for 15 min (after centrifuged 1 mL of sample was again centrifuged at 17,000 g for 15 min) and filtered through 0.45 µm membrane filter. Instrumental conditions during free amino acid analysis are as follows: S433, auto analyzer, Cation separation column (LCAK07/li), 4.6×150 mm; buffer change (A, pH 2.90; B, pH 4.20; C, pH 8.00); (Lithium citrate buffer solution), buffer flow rate: 0.45 mL/min, ninhydrin flow rate: 0.25 mL/min, Column temp.: 37°C.

Statistical analysis

Statistical analysis was done by GLM procedure using (SAS, 2003). A significance test between the results was used by Tukey's multiple test method ($p < 0.05$). The results of the analysis are presented as standard errors of the mean and the least-square means.

Results and Discussion

Carnosine and anserine

The carnosine and anserine contents of HH, broiler, 2A, 2C, and 2D from male breast meat are presented in Fig. 1. The concentration of carnosine in HH, broiler, 2A, 2C, and 2D were 511.04 (mg/100 g), 257.94 (mg/100 g), 287.39 (mg/100 g), and 359.31 (mg/100 g) respectively. The carnosine content was highest in HH and lowest in broiler ($p < 0.05$). The concentration of carnosine was more dominant in native chickens than broiler which was an agreement with the previous study stated by (Intarapichet and Maikhunthod, 2005). It was found that the new native chicken strains, 2A, 2C, and 2D contained more carnosine content than broiler (Fig. 1). Comparing of native chicken strains with broiler, these results were consistent with those reported by (Jung et al., 2013) in Korean native chicken. This is due to different composition of muscle fiber type I (slow-twitch oxidative red fiber), IIA (fast-twitch oxidative-glycolytic white fiber), IIB (fast-twitch glycolytic white fiber) in meat (Cornet and Bousset, 1999). The inconsistent of carnosine in different chickens may be due to genotypic trait that could be different among traits such as enzyme and transporters agreed by (Everaert et al., 2013). Higher concentration of carnosine has been found with the higher concentration of white muscle rather than the tissue of red muscle that continue their energy-rich phosphate ester in a condition of anaerobic means (Boldyrev et al., 2004). There for further studies are needed to elucidate the biological significance of carnosine among the native chicken strains. The anserine content of HH, broiler, 2A, 2C, and 2D from male breast meat were 1,525.80 (mg/100 g), 660.38 (mg/100 g), 1,286.34 (mg/100 g), 962.70 (mg/100 g), and 1,059.76 (mg/100 g) respectively. It was found that HH contained significantly ($p < 0.05$) higher




Fig. 1. Carnosine, anserine, and reducing sugar contents (mg/100 g) of the breast meats from HH, broiler, 2A, 2C, and 2D. ^{a-c} Different small letter indicate the statistically significant ($p < 0.05$) in breast meat. HH, commercial native chicken (HanHyup No. 3); 2A, 2C, and 2D, new native chicken strains; broiler, commercial broiler.

anserine content among all groups of chickens. The new native chicken strains, 2A, 2C, and 2D contained higher amount of anserine than broiler. Anserine is the predominating histidine dipeptide in poultry meat, whereas carnosine for beef and pork (Abe, 1995). Anserine (1-methylated carnosine) derived from carnosine mostly abundant in non-mammalian species, such as poultry agreed with the results conducted by (Abe, 1995). Anserine is produced by catalyzed of carnosine as it is the derived product of carnosine (Drozak et al., 2013). The effect of meat type on anserine content corresponded with that on carnosine content. Native chickens contained high anserine than broiler that was due to the activity of anserine act as a physico-chemical buffer against proton production by anaerobic glycolysis in different muscle traits (Dunnnett and Harris, 1995).

Reducing sugar

The reducing sugar contents of HH, broiler, 2A, 2C, and 2D from male breast meat are shown in (Fig. 1). The reducing sugar concentrations of HH, broiler, 2A, 2C, and 2D were 78.32 (mg/100 g), 44.87 (mg/100 g), 87.55 (mg/100 g), 93.47 (mg/100 g), and 75.01 (mg/100 g) respectively. Among the groups, it was found that new native chicken strain, 2C contained significantly higher reducing sugar than broiler ($p < 0.05$). And, between native chickens and broiler, the broiler had lower reducing sugar in breast meat than other native chickens. In general, the native chicken contained higher reducing sugar than

broiler supported by (Jung et al., 2013). The reducing sugars responsible for some bothy and aroma flavor in chicken meat (Sasaki et al., 2007). (Mottram, 1998) reported that free sugars, sugar phosphates, nucleotide bound sugars, free amino acids, peptides, nucleotides and other nitrogenous components, such as thiamine as the main water-soluble flavor precursors varies according to genetic makeup and muscle structure. The reaction of cysteine and sugar can lead to characteristic meat flavour specially for chicken (Varavinit et al., 2000). During heating of breast meat, ribose which is a reducing sugar in meat associated with ribonucleotides in the muscle produce the flavor (Mottram, 1998).

Vitamins and cholesterol

The vitamins (α -tocopherol, γ -tocopherol, α -tocotireinol, Vitamin A, and Vitamin B₁₂) contents of HH, broiler, 2A, 2C, and 2D from male breast fillet are presented in Table 1. The new native chicken strains, 2A, 2C, and 2D contained higher levels of α -tocopherol than HH and broiler but among the groups α -tocopherol content was highest in 2C which was 0.26 (mg/100 g) and showed a significant difference among groups ($p < 0.05$). Vitamin A was higher in 2A among the groups but did not show any significant difference ($p > 0.05$). New native strains, 2C, 2D, and HH showed the lower contents of vitamin A than broiler that agreed with study conducted by (Lee et al., 2015). Cholesterol contents of HH, broiler, 2A, 2C, and 2D were 51.26 (mg/100 g), 63.17 (mg/100 g), 54.43 (mg/100 g), 49.04 (mg/100 g), and 49.51 (mg/100 g) respectively. It is showed that (Table 1) the highest cholesterol content was found in broiler which was 63.17 (mg/100 g) and higher than all native chickens that was agreed by (Jaturasitha et al., 2008). But among new native chicken strains, 2C and 2D implied the lower inflate of cholesterol than HH. It is, however, the cholesterol level varies according to contents of body fat, skin fat, and bone marrow (Demos and Mandigo, 1995).

Minerals

Mineral contents of HH, broiler, 2A, 2C, and 2D from male breast fillet are given in Table 2. The predominant minerals, phosphorous was significantly higher in broiler than that of all treatment groups. For magnesium, broiler also contained more contents than all groups. But in 2A and 2C had higher concentration of magnesium than HH (Table 2). New native chicken strains, 2A, 2C, and 2D had higher sodium contents than HH. Although P, Mg, and Na were higher in broiler compared with other groups but these minerals have no health effects in human body. Lower calcium content in each treatment indicates the less bone ion meat. The micronutrients investigated in this study were zinc, iron, manganese and copper. 2A had higher zinc contents than HH, 2C, and 2D. Higher iron found in HH compared with all groups. Meat with zinc and iron has a great

Table 1. Vitamins and cholesterol contents of the breast meats from HH, broiler, 2A, 2C, and 2D

Items	HH	Broiler	2A	2C	2D	SEM ¹⁾
α -Tocopherol (mg/100 g)	0.23 ^d	0.17 ^e	0.24 ^c	0.26 ^a	0.25 ^b	0.00
γ -Tocopherol (mg/100 g)	0.04 ^b	0.05 ^a	0.04 ^b	0.05 ^a	0.04 ^a	0.00
α -Tocotireinol (mg/100 g)	0.012 ^a	0.010 ^{ab}	0.012 ^a	0.012 ^a	0.008 ^b	0.00
Vitamin A (μ g/100 g)	4.94	5.65	5.88	5.43	5.25	0.35
Vitamin B ₁₂ (mg/100 g)	0.30 ^a	0.28 ^{ab}	0.27 ^{ab}	0.24 ^b	0.28 ^{ab}	0.01
Cholesterol (mg/100 g)	51.26 ^c	63.17 ^a	54.43 ^b	49.04 ^c	49.51 ^c	0.61

¹⁾ Standard error of the means (n=40).

^{a-c} Different letters within the same column differ significantly ($p < 0.05$).

HH, commercial native chicken (HanHyup No. 3); 2A, 2C, and 2D, new native chicken strains; Broiler, commercial broiler.

Table 2. Mineral contents of the breast meats from HH, broiler, 2A, 2C, and 2D

Minerals	HH	Broiler	2A	2C	2D	SEM ¹⁾
Ca (%)	0.01	0.01	0.01	0.01	0.01	0.00
P (%)	0.21 ^{bc}	0.23 ^a	0.20 ^c	0.21 ^b	0.20 ^c	0.00
Mg (ppm)	318.37 ^b	335.23 ^a	318.23 ^b	321.85 ^b	315.52 ^b	1.73
Na (ppm)	296.34 ^c	409.22 ^a	314.99 ^b	302.21 ^c	299.10 ^c	1.61
Zn (ppm)	5.55 ^b	7.18 ^a	5.68 ^b	5.31 ^b	5.35 ^b	0.22
Fe (ppm)	9.37	8.85	7.70	8.00	5.00	1.12
Mn (ppm)	0.42	0.23	0.40	0.25	0.07	0.10
Cu (ppm)	0.34 ^a	0.26 ^b	0.35 ^a	0.28 ^b	0.30 ^{ab}	0.01

¹⁾ Standard error of the means (n=40).

^{a-c} Different letters within the same column differ significantly (p<0.05).

HH, commercial native chicken (HanHyup No. 3); 2A, 2C, and 2D, new native chicken strains; Broiler, commercial broiler.

importance to reduce the incidence of some diseases and being accumulated as a part of beneficial effect on health (Lombardi-Boccia et al., 2005). Copper content was higher in 2A from all treatment groups. The contents of mineral can be influenced by many factors such as species, breeding condition, age, and supplementation with various ingredients (Park and Kim, 2001).

Free amino acids

Free amino acid concentrations of HH, broiler, 2A, 2C, and 2D from male breast fillet are depicted in Table 3. In the fraction of essential amino acids, the predominant amino acids were the tryptophan and threonine whereas alanine was superior found in the non-essential fraction, followed by serine while lowest values were reported for arginine among treatments. Umami or savory amino acid, glutamic acid content in 2A was highest among the all groups (p<0.05). Glutamic acid which is one of the most important amino acid in chicken meat regarding of enhancing the flavor of meat either combination with other test-associated compounds or its by self (Kurihara, 1987). Glutamic acid in native chicken was higher than broiler that is agreed with the study conducted by (Ahn and Park, 2002). The flavor related amino acids, valine, isoleucine, leucine, phenylalanine, arginine, and proline were significantly higher in 2A among the groups (p<0.05). Subsequently HH contained more flavor producing amino acid than broiler (Table 3). The Korean native chickens are superior in flavor than broiler supported by (Choe et al., 2010). It is however, tasty amino acids, asparagine, threonine, serine, glutamic acid, glycine, and alanine contents were significantly higher in 2A than HH and broiler (p<0.05). The more functional compound, taurine contents of HH, broiler, 2A, 2C, and 2D were 5.76 (ppm), 4.18 (ppm), 18.66 (ppm), 4.71 (ppm), and 6.18 (ppm) respectively (Table 3), while 2A contained significantly higher taurine content than other groups (p<0.05) which are not association with taste or odor of meat (Batzer et al., 1960). The native chickens compared with broiler had higher taurine content supported by (Choe et al., 2010). Taurine influences many functions like as bile salt formation, stabilization of membrane, calcium homeostasis, and growth moderator (Redmond et al., 1998). And, moreover, broiler had higher cysteine which is sulfur containing amino acid but lower content of methionine than other groups. The amino acid which performs the immune functions in the human body was significantly higher in 2A among the groups (p<0.05).

Conclusion

In general, the new native chicken strains have more carnosine, anserine, and reducing sugar than broiler. Due to higher

Table 3. Free amino acid concentration (ppm) of the breast meats from HH, broiler, 2A, 2C, and 2D

Free amino acids	HH	Broiler	2A	2C	2D	SEM ¹⁾
Taurine	5.76 ^b	4.18 ^b	18.66 ^a	4.71 ^b	6.18 ^b	0.50
Aspartic acid	11.83 ^b	5.71 ^c	22.45 ^a	6.95 ^c	11.04 ^b	1.18
Threonine	16.18 ^a	0.18 ^b	18.64 ^a	12.59 ^a	14.27 ^a	1.71
Serine	21.09 ^b	15.75 ^b	29.28 ^a	17.36 ^b	18.49 ^b	2.34
Asparagine	1.55 ^d	17.93 ^b	33.17 ^a	8.92 ^c	21.75 ^b	1.87
Glutamic acid	21.66 ^b	9.40 ^c	28.30 ^a	8.95 ^c	20.73 ^b	1.72
Glycine	22.24 ^b	15.99 ^b	47.14 ^a	16.04 ^b	18.40 ^b	2.43
Alanine	28.24 ^b	19.13 ^b	42.54 ^a	20.34 ^b	25.46 ^b	2.43
Valine	10.57 ^b	13.95 ^a	7.32 ^{bc}	5.22 ^c	6.94 ^{bc}	1.00
Cystine	0.85 ^b	6.10 ^a	0.45 ^c	0.19 ^d	0.15 ^d	0.07
Methionine	8.31 ^a	4.88 ^b	7.19 ^{ab}	6.63 ^{ab}	7.84 ^a	0.63
Isoleucine	10.57 ^b	13.95 ^a	7.32 ^{bc}	5.22 ^{bc}	6.94 ^c	1.00
Leucine	21.84 ^a	7.03 ^c	19.59 ^{ab}	14.39 ^b	17.79 ^{bc}	1.35
Tyrosin	9.99 ^a	5.68 ^b	10.27 ^a	7.60 ^{ab}	8.51 ^{ab}	0.86
Phenylalanine	8.93 ^a	1.32 ^c	8.19 ^{ab}	5.81 ^b	7.11 ^{ab}	0.64
Tryptophan	21.12 ^a	12.48 ^{bc}	7.64 ^c	16.67 ^{ab}	15.78 ^{ab}	1.84
Lysine	14.45 ^a	7.20 ^b	14.73 ^a	8.44 ^b	9.55 ^b	0.82
Arginine	0.46 ^b	0.10 ^b	1.07 ^a	0.29 ^b	0.81 ^a	0.11
Tasty AA ²⁾	110.95 ^b	78.39 ^b	199.07 ^a	84.20 ^b	119.09 ^b	10.36
FRAA ³⁾	85.27 ^b	56.44 ^c	141.95 ^a	52.76 ^c	76.59 ^{bc}	7.14

¹⁾ Standard error of the means (n=40).

²⁾ Tasty AA: Sum of asparagine, threonine, serine, glutamic acid, glycine, and alanine.

³⁾ FRAA (Flavor-Related AA): Sum of valine, isoleucine, leucine, phenylalanine, arginine, and proline.

^{a-c} Different letters within the same column differ significantly (p<0.05).

HH, commercial native chicken (HanHyup No. 3); 2A, 2C, and 2D, new native chicken strains; Broiler, commercial broiler.

tocopherol and lower cholesterol values, some native chicken strains may be accepted to the consumers with health benefits aspects. The new native chicken strain have more umami taste compare to broiler or HH. Tasty and flavor related amino acids of new native chicken strains are better than broiler or HH. Comparing with HH or broiler, the new native chicken strains have some distinct superior characters that lead to select in poultry breeding sector. So, this work may be regarded as benchmark for poultry breeder to take into consideration in breeding policy to get more nutritional meat with desired quality.

Conflict of Interest

The authors declare no potential conflict of interest.

Acknowledgements

This study was supported by Golden Seed Project (No. 213010-05-1-SB420, PJ01282202), Ministry of Agriculture, Food and Rural Affairs, Ministry of Oceans and Fisheries, Rural Development Administration, Korea.

Author Contributions

Conceptualization: Nam KC. Data curation: Park JY, Lee SY, Jung S. Formal analysis: Park JY, Lee SY. Methodology: Park JY, Lee SY, Jung S. Validation: Jo C. Writing - original draft: Ali M. Writing - review & editing: Ali M, Lee SY, Park JY, Jung S, Jo C, Nam KC.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Abe H, Okuma E. 1995. Discrimination of meat species in processed meat-products based on the ratio of histidine dipeptides. *J Jpn Soc Food Sci* 42:827-834.
- Ahn DH, Park SY. 2002. Studies on components related to taste such as free amino acids and nucleotides in Korean native chicken meat. *J Korean Soc Food Sci Nutr* 31:547-552.
- Aliani M, Farmer LJ. 2005. Precursors of chicken flavor. I. Determination of some flavor precursors in chicken muscle. *J Agric Food Chem* 53:6067-6072.
- Baguet A, Koppo K, Pottier A, Derave W. 2010. Beta-alanine supplementation reduces acidosis but not oxygen uptake response during high-intensity cycling exercise. *Eur J Appl Physiol* 108:495-503.
- Batzer OF, Santoro AT, Tan MC, Landmann WA, Schweigert BS. 1960. Meat flavor chemistry, precursors of beef flavor. *J Agric Food Chem* 8:498-501.
- Boldyrev A, Bulygina E, Leinsoo T, Petrushanko I, Tsubone S, Abe H. 2004. Protection of neuronal cells against reactive oxygen species by carnosine and related compounds. *Comp Biochem Physiol B Biochem Mol Biol* 137:81-88.
- Choe JH, Nam KC, Jung S, Kim BN, Yun HJ, Jo CR. 2010. Differences in the quality characteristics between commercial Korean native chickens and broilers. *Korean J Food Sci Anim Resour* 30:13-19.
- Chowdhury SR, Chowdhury SD, Smith TK. 2002. Effects of dietary garlic on cholesterol metabolism in laying hens. *Poult Sci* 81:1856-1862.
- Cornet M, Bousset J. 1999. Free amino acids and dipeptides in porcine muscles: Differences between 'red' and 'white' muscles. *Meat Sci* 51:215-219.
- Demos BP, Mandigo RW. 1995. Composition and chemistry of mechanically recovered beef neck-bone lean. *J Food Sci* 60:576-579.
- Drozak J, Chrobok L, Poleszak O, Jagielski AK, Derlacz R. 2013. Molecular identification of carnosine n-methyltransferase as chicken histamine n-methyltransferase-like protein (hnmt-like). *PLOS ONE* 8:e64805.
- Dunnett M, Harris RC. 1995. Carnosine and taurine contents of type i, iia and iib fibres in the middle gluteal muscle. *Equine Vet J* 27:214-217.
- Everaert I, De Naeyer H, Taes Y, Derave W. 2013. Gene expression of carnosine-related enzymes and transporters in skeletal muscle. *Eur J Appl Physiol* 113:1169-1179.
- Food and Drug Administration. Korean Food Standards Codex. Korea Food and Drug Administration, Cheongju, Korea.
- Hipkiss AR. 2009. Carnosine, diabetes and alzheimer's disease. *Expert Rev Neurothe* 9:583-585.

- Intarapichet KO, Maikhunthod B. 2005. Genotype and gender differences in carnosine extracts and antioxidant activities of chicken breast and thigh meats. *Meat Sci* 71:634-642.
- Iwaniak A, Dziuba B. 2009. Motifs with potential physiological activity in food proteins: Biopep database. *Acta Sci Pol Technol Aliment* 8:59-85.
- Jaturasitha S, Srikanchai T, Kreuzer M, Wicke M. 2008. Differences in carcass and meat characteristics between chicken indigenous to northern Thailand (Black-boned and Thai native) and imported extensive breeds (Bresse and Rhode Island Red). *Poult Sci* 87:160-169.
- Jung S, Bae YS, Kim HJ, Jayasena DD, Lee JH, Park HB, Heo KN, Jo C. 2013. Carnosine, anserine, creatine, and inosine 5'-monophosphate contents in breast and thigh meats from 5 lines of Korean native chicken. *Poult Sci* 92:3275-3282.
- Kurihara K. 1987. Recent progress in the taste receptor. In *Umami: A basic taste*. Marcel Dekker, New York, NY, USA.
- Lee JH, Lee HN, Shin JA, Chun JY, Lee J, Lee KT. 2015. Content of fat-soluble nutrients (cholesterol, retinol, and α -tocopherol) in different parts of poultry meats according to cooking method. *J Korean Soc Food Sci Nutr* 44:234-241.
- Lombardi-Boccia G, Lanzi S, Aguzzi A. 2005. Aspects of meat quality: Trace elements and b vitamins in raw and cooked meats. *J Food Compos Anal* 18:39-46.
- Meinert L, Schafer A, Bjerregaard C, Aaslyng MD, Bredie WL. 2009. Comparison of glucose, glucose 6-phosphate, ribose, and mannose as flavour precursors in pork; the effect of monosaccharide addition on flavour generation. *Meat Sci* 81:419-425.
- Mora L, Sentandreu MA, Toldra F. 2007. Hydrophilic chromatographic determination of carnosine, anserine, balenine, creatine, and creatinine. *J Agric Food Chem* 55:4664-4669.
- Morrissey PA, Sheehy PJ, Galvin K, Kerry JP, Buckley DJ. 1998. Lipid stability in meat and meat products. *Meat Sci* 49S1:S73-S86.
- Mottram D, Maarse H. 1991. Volatile compounds in foods and beverages. Marcel Dekker, New York, NY, USA. pp 107-177.
- Mottram DS. 1998. Flavour formation in meat and meat products: A review. *Food Chem* 62:415-424.
- Mun GJ, Song W, Park SH, Jeong BG, Jung GR, Choi KS, Kim JH, Choi Y, Chun J. 2017. Analysis of vitamin B₁₂ in fresh cuts of Korean pork for update of national standard food composition table. *Korean J Food Preserv* 24:983-991.
- Nagai H, Harada M, Nakagawa M, Tanaka T, Gunadi B, Justinus Setiabudi ML, Uktolseja JL, Miyata Y. 1996. Effects of chicken extract on the recovery from fatigue caused by mental workload. *Appl Hum Sci* 15:281-286.
- Ouali A, Herrera-Mendez CH, Coulis G, Becila S, Boudjellal A, Aubry L, Sentandreu MA. 2006. Revisiting the conversion of muscle into meat and the underlying mechanisms. *Meat Sci* 74:44-58.
- Park C, Kim Y. 2001. Effect of additions of supplemental activated carbon on the fatty acid, meat color and minerals of chicken meat. *Korean J Food Sci Anim Resour* 21:285-291.
- Park HB, Seo DW, Choi NR, Choi JS, Heo KN, Kang BS, Jo C, Lee JH. 2012. Estimation of genetic parameters for serum clinical-chemical traits in Korean native chickens. *Korean J Poult Sci* 39:279-282.
- Pereira PM, Vicente AF. 2013. Meat nutritional composition and nutritive role in the human diet. *Meat Sci* 93:586-592.
- Redmond HP, Stapleton PP, Neary P, Bouchier-Hayes D. 1998. Immunonutrition: The role of taurine. *Nutrition* 14:599-604.
- Ribeiro DM, Mourato MP, Almeida AM. 2019. Assessing mineral status in edible tissues of domestic and game animals: A review with a special emphasis in tropical regions. *Trop Anim Health Prod* 51:1-14.
- SAS. 2003. SAS program. Version 9.4, SAS Institute Inc., Cary, NC, USA.
- Sasaki K, Motoyama M, Mitsumoto M. 2007. Changes in the amounts of water-soluble umami-related substances in porcine

- longissimus* and *biceps femoris* muscles during moist heat cooking. *Meat Sci* 77:167-172.
- Schmid A. 2009. Bioactive substances in meat and meat products. *Fleischwirtschaft* 89:83-90.
- Simopoulos AP. 2004. Omega-6/omega-3 essential fatty acid ratio and chronic diseases. *Food Rev Int* 20:77-90.
- The Korea Society of Food Science and Nutrition. 2000. Handbook of experiments in food science and nutrition. Hyoil Publishing Co., Seoul, Korea.
- Thomas JB, Kline MC, Gill LM, Yen JH, Duewer DL, Sniegoski LT, Sharpless KE. 2001. Preparation and value assignment of standard reference material 968c fat-soluble vitamins, carotenoids, and cholesterol in human serum. *Clin Chim Acta* 305:141-155.
- Varavinit S, Shobsngob S, Bhidyachakorawat M, Suphantharika M. 2000. Production of meat-like flavor. *Sci Asia* 26:219-224.
- WHO. 2003. Diet, nutrition, and the prevention of chronic diseases. World Health Organization, Geneva, Switzerland. p 149.
- Yamaguchi S. 1991. Roles and efficacy of sensory evaluation in studies of taste. *J Jpn Soc Food Sci* 38:972-978.