

Comparison of pH, Water Holding Capacity and Color among Meats from Korean Native Chickens

Samooel Jung¹, Hyun Joo Kim², Hyun Jung Lee², Dong Won Seo¹, Jun Heon Lee¹, Hee Bok Park¹,
Cheorun Jo² and Ki Chang Nam^{3,†}

¹Department of Animal Science and Biotechnology, Chungnam National University, Daejeon 305-764, Korea

²Department of Agricultural Biotechnology, Center for Food and Bioconvergence, and Research Institute of Agriculture and Life Sciences, Seoul National University, Seoul 151-921, Korea

³Department of Animal Science and Technology, Suncheon National University, Suncheon 540-742, Korea

ABSTRACT This study investigated the initial pHs measured 15 min post-mortem, the ultimate pHs, the water holding capacities (WHCs), and the colors of meats obtained from five lines of Korean native chicken (KNC) for the development of a breed for high-quality meat. Additionally, the effect of sex was examined. In total, 595 F1 progeny (Black [B], Grey-Brown [G], Red-Brown [R], White [W] and Yellow-Brown [Y]) from 70 full-sib families were used. The chickens were slaughtered at 20 wk of age, and the measurement traits of all breast and thigh meats from the 595 chickens were analyzed. The initial pHs 15 min post-slaughter of the breast and thigh meats were affected by the line and sex of the Korean native chickens ($P<0.05$). However, there were no line or sex effects on the ultimate pHs and WHCs of the breast and thigh meats, with the exception that the ultimate pH of the thigh meat exhibited a line effect. With the exception of the L^* value of the breast meat, the meat color was significantly affected by line and sex ($P<0.05$). The ultimate pHs exhibited consistently negative correlations with the L^* values and positive correlations with the a^* values of the breast and thigh meats. Based on these results, we concluded that the line W male chickens and the line G and W female chickens may be good candidates for the development of a breed for high-quality meat because these lines exhibited the properties of high initial pHs or/and ultimate pHs.

(Key words: Korean native chicken breeds, pH, water holding capacity, color)

INTRODUCTION

Producers of chicken meat have attempted to find a way to produce high-quality meat to satisfy the increased consumer demand. There is a growing interest in the use of native chicken breeds for the production of chicken meat in many countries because native chicken meats have properties such as high protein content, low fat content, unique flavor and texture compared with those of broiler chicken meat (Fanatico et al., 2007; Jaturasitha et al., 2008; Jayasena et al., 2013; Jung et al., 2013). In addition, the conservation of native chicken breeds is considered important because the increasing use of broiler chickens with rapid growth properties has resulted in the loss of genetic diversity among chicken breeds (Zanetti et al., 2010). For these reasons, the development of a new meat breed by using native chickens is required, based on suitable selection strategies for the production of high-quality meat (Jung et al., 2013).

Meat quality is strongly associated with the decline of muscle pH post-mortem. Rapid decline of muscle pH and low ultimate pH values result in pale, soft and exudative (PSE) meat that has a pale appearance, soft texture and exudative traits owing to decreased water holding capacity (WHC); denaturation of meat chromoproteins, especially myoglobin; and alteration of muscle fiber structure (Owens et al., 2000; Swatland, 2008). In contrast, high ultimate pH generates dark, firm and dry (DFD) meat with a dark appearance, firm texture, and dry traits, in addition to poor storage stability (Allen et al., 1997). Although PSE and DFD meat are primary concerns for pork and beef, the incidence of PSE meat has been reported for chicken meat (Garcia et al., 2010; Barbut et al., 2008). Therefore, regulating the rate and extent of muscle pH decline after slaughter is crucial to the production of chicken meat. Barbut et al. (2008) emphasized the importance of ultimate pH, rate of pH decline, WHC and color of meat in the selection process.

[†] To whom correspondence should be addressed : kichang@sncu.kr

Recently, government agencies in Korea have been engaged in a project to develop a new meat breed of chicken based on Korean native chickens (KNCs), to conserve native chicken breeds while simultaneously satisfying consumer demands. Five KNC lines (i.e., Black, Grey-Brown, Red-Brown, White, and Yellow-Brown) have been proposed as candidates for the selection (Jung et al., 2013). The present study was conducted to investigate the pH, WHC and color of breast and thigh meat obtained from the five KNC lines to obtain useful information for devising selection strategies.

MATERIALS AND METHOD

1. Animals

A two-generation resource pedigree using five lines of KNC was established and managed in this study. Within each line which was established in National Institute of Animal Science (NIAS) of Korea (Cheonan, Korea), three sires were mated with 14 to 15 dams to produce F₁ chicks. In total, 595 F₁ progeny (Black [B]: 90 [male: 45, female: 45], Grey-Brown [G]: 110 [male: 52, female: 58], Red-Brown [R]: 136 [male: 68, female: 68], White [W]: 126 [male: 63, female: 63], Yellow-Brown [Y]: 133 [male: 62, female: 71]) from 70 full-sib families were used in this study. Chickens were raised at NIAS (Cheonan, Korea) and fed *ad libitum* commercial-formula feed containing 18.2% concentrated protein and 2859 kcal/kg metabolizable energy. Chicken care facilities and the procedures were performed met or exceeded the standards established by the Committee for Accreditation of Laboratory Animal Care at NIAS in Korea. This study was also conducted in accordance with "The Guide for the Care and Use of Laboratory Animals" published by the institutional Animal Care and Use Committee of NIAS, Korea. Chickens were slaughtered after 4 h of feed withdrawal at 20 wk of age using conventional neck cuts and bleeding for 2 min; the feathers were then removed and the chickens were eviscerated. The carcasses were vacuum-packed after chilling in ice-cold water and stored in refrigerator at 4°C for 24 h. And then vacuum-packed carcasses were frozen in a freezer at -20°C until analysis.

2. Sample Preparation

Before analysis, the frozen carcasses were thawed in a refrigerator at 4°C for 24 h. Breast and thigh muscles were dissected from each thawed carcass. Then, the right and left breast and thigh muscles from each chicken were minced separately using a food mixer (CH180A, Kenwood Ltd, Hampshire, UK) for 30 sec. Minced meat samples were used for analysis.

3. Measurement of pH

After slaughter, the pH values at 15 min post-slaughter were measured by inserting the probe into the respective breast and thigh muscles of fresh carcass with portable pH meter (SG2-SevenGo, Mettler-Toledo Inti Inc., Schwerzenbach, Switzerland). To measure ultimate pH values, minced meat (1 g) of respective breast and thigh muscles from frozen-thawed carcasses were homogenized with 9 mL of distilled water using a homogenizer (T25 basic, IKA GmbH & Co. KG, Germany) at $1,130 \times g$ for 30 sec. The homogenates were filtered by filter paper (No. 4, Whatman Ltd. Kent, UK) after centrifugation at $2,090 \times g$ for 15 min (Union 32R, Hanil Co., Ltd., Incheon, Korea). The pH of filtrate was measured using a pH meter (SevenEasy, Mettler-Toledo Inti Inc., Schwerzenbach, Switzerland).

4. Measurement of Water Holding Capacity

One gram of the minced meat sample was placed on a round filter paper (No.4, Whatman Ltd. Kent,UK). The filter paper with meat was placed into centrifuge tube and the tube was centrifuged (CR 20B2, Hitachi Koki Co., Ltd. Fukuoka, Japan) at $6,710 \times g$ for 10 min. The released water absorbed into the filter paper was weighed and calculated as a percentage of the initial moisture of meat. Moisture content of meat was determined by drying 3 g of samples placed in aluminum dishes for 15 h at 104°C.

5. Measurement of Instrumental Color

The CIE lightness (L^*), redness (a^*) and yellowness (b^*) of the minced meat were measured with a colorimeter (CR-300, Minolta Inc., Tokyo, Japan). This instrument was calibrated with a black and a white reference tile. Measurements were taken perpendicularly to the sample in quartz cell (3 cm \times 1.5 cm) at two different locations per sample. The result was analyzed using Spectra Magic Software (Minolta, Japan)

automatically.

6. Statistical Analysis

All data (pooled data) were analyzed by multifactorial analysis of variance using the general linear model to investigate the effect of meat type (breast and thigh), gender (male and female), and line (five lines of KNC). After grouping the data by each meat type with each gender, the data were analyzed by the general linear model to confirm the line effect in each meat type with each gender. Tukey's multiple range test was used to compare significant differences between least square mean values ($P<0.05$). Least square mean values and standard error of the least square means (SEM) are reported. Additionally, Pearson's correlation coefficients were calculated ($P<0.05$). SAS software (version 9.3, SAS Institute Inc., Cary, NC, USA) was used for all statistical analyses.

RESULTS AND DISCUSSION

1. pH and Water Holding Capacity

Energy production for muscle contraction in postmortem muscle mainly depends on anaerobic glycolytic metabolism that leads to accumulation of lactic acid in the muscle before rigor mortis (Dransfield and Sosnicki, 1999). Therefore, post-mortem muscle undergoes a decrease in pH, and the degree

and rate of pH decline can be affected by not only the environment but also genetic properties (Debut et al., 2003). In the present study, the five KNC lines were raised and slaughtered under the same environment to exclude the effect of environment on muscle pH.

The initial pH at 15 min post-slaughter and the ultimate pH of breast meat obtained from lines B, G, R, W and Y of KNC are presented in Table 1. The initial pH of breast meat from male chickens was significantly higher in line W than in lines B and Y ($P<0.05$), whereas no significant difference in initial pH was observed in breast meat from female chickens. The low pH value at 15 min post-slaughter in chicken meat indicates a rapid decline in pH. However, the ultimate pH of breast meat from both male and female chickens did not significantly differ among the five KNC lines.

No significant difference in WHC was observed for breast meat from male or female chickens among the five KNC lines, although the initial pH of breast meat from male chickens was significantly different. The WHC of meat reflects the affinity between myofibrillar protein and water molecules, and the space to retain water within the myofibril (Huff-Lonerger and Lonergan, 2005). The decline of pH near the isoelectric point of myofibrillar protein results in low WHC owing to reduction of the net charge of myofibrillar protein, thereby decreasing both water affinity and space (Huff-Lonerger and Lonergan,

Table 1. pH and water holding capacity (%) of the breast meats from five lines of Korean native chicken

| Line ¹ | pH _{15min} | | pH _{ultimate} | | WHC | |
|-------------------|---------------------|--------|------------------------|--------|--------|--------|
| | Male | Female | Male | Female | Male | Female |
| B | 6.08 ^c | 6.18 | 5.77 | 5.78 | 64.40 | 63.71 |
| G | 6.23 ^{ab} | 6.25 | 5.74 | 5.76 | 63.07 | 64.36 |
| R | 6.21 ^{ab} | 6.31 | 5.81 | 5.78 | 64.87 | 62.06 |
| W | 6.28 ^a | 6.27 | 5.81 | 5.82 | 61.86 | 64.12 |
| Y | 6.17 ^{bc} | 6.24 | 5.83 | 5.79 | 64.44 | 64.08 |
| SEM | 0.031 | 0.038 | 0.027 | 0.031 | 1.121 | 0.929 |
| | Line | Gender | Line | Gender | Line | Gender |
| <i>P</i> -value | 0.0010 | 0.0200 | 0.0503 | 0.5190 | 0.7251 | 0.8346 |
| <i>F</i> -value | 6.64 | 6.31 | 2.32 | 0.42 | 0.51 | 0.04 |

^{a~c} Different letters among breeds differ significantly ($P<0.05$).

¹ B=Black, G=Grey-Brown, R=Red-Brown, W=White and Y=Yellow-Brown.

2005). Debut et al. (2003) reported that both low ultimate pH and accelerated rate of pH decline (low pH at 15 min post-slaughter) could lead to poor WHC of chicken meat. Le Bihan-Duval et al. (2008) found that ultimate pH had higher correlation with WHC than did initial pH (measured at 15 min post-slaughter) in broiler chickens. Therefore, no significant difference in WHC of breast meat among the five KNC lines might be consequence of no significant difference of ultimate pH of the breast meats among the five KNC lines.

Female chickens of lines G and W had significantly higher initial and ultimate pH values in the thigh meat than did females of line R ($P<0.05$), whereas no significant difference in initial and ultimate pH values was observed in male chickens among the five KNC lines. However, the WHC of thigh meat did not differ among the five KNC lines for both male and female chickens. The inconsistency of the relationship between ultimate pH and WHC in meat was previously reported in different chicken breeds. Musa et al. (2006) found a significant difference of up to 0.04 pH units in ultimate pH, with no difference in WHC in chicken breast meat. Jung et al. (2011) found no significant difference in WHC in chicken thigh meat, although a significant difference of 0.17 pH units was observed for ultimate pH. However, a significant difference of 0.20 pH units in ultimate pH resulted in a difference in WHC in chicken thigh meat (Debut et al., 2003). In the present study,

the ultimate pH in thigh meat obtained from male and female KNC ranged from 6.11 to 6.23. Therefore, the difference in ultimate pH, if any, was likely too small to influence the WHC of thigh meat obtained from both male and female KNC.

In the present study, the effects of sex (male and female) on the pH and WHC of meat obtained from the five KNC lines, from pooled data, were investigated, in addition to investigating the effect of the chicken line. The initial pH of breast meat and thigh meat obtained from male and female chickens was 6.20 and 6.25 and 6.51 and 6.55, respectively, in the KNC lines (data not shown, $P<0.05$). However, no effects of sex on the ultimate pH and WHC of both breast and thigh meat from the KNC lines was observed. These results agree with those of a previous study. Musa et al. (2006) reported that there was no effect of sex on the ultimate pH and WHC of chicken meat.

2. Instrumental Meat Color

The instrumental color (lightness L^* , redness a^* and yellowness b^*) of breast meat obtained from the five KNC lines is presented in Table 3. From the pooled data, an effect of chicken line was observed on the a^* and b^* values. The a^* value of breast meat from female chickens was significantly higher in line B than in line Y ($P<0.05$). However, there was no significant difference in the a^* value for breast meat obtained from male chickens of the five KNC lines. The breast meat

Table 2. pH and water holding capacity (%) of the thigh meats from five lines of Korean native chicken

| Line ¹ | pH _{15min} | | pH _{ultimate} | | WHC | |
|-------------------|---------------------|--------------------|------------------------|--------------------|--------|--------|
| | Male | Female | Male | Female | Male | Female |
| B | 6.50 | 6.48 ^b | 6.15 | 6.16 ^{ab} | 63.66 | 60.49 |
| G | 6.55 | 6.60 ^a | 6.18 | 6.23 ^a | 60.52 | 61.40 |
| R | 6.52 | 6.50 ^b | 6.18 | 6.11 ^b | 61.44 | 61.35 |
| W | 6.50 | 6.59 ^a | 6.18 | 6.21 ^a | 60.55 | 61.26 |
| Y | 6.51 | 6.55 ^{ab} | 6.22 | 6.22 ^a | 61.99 | 62.22 |
| SEM | 0.031 | 0.028 | 0.020 | 0.025 | 0.816 | 0.789 |
| | Line | Gender | Line | Gender | Line | Gender |
| <i>P</i> -value | 0.0456 | 0.0195 | 0.0006 | 0.5707 | 0.4450 | 0.9752 |
| <i>F</i> -value | 2.45 | 5.49 | 4.96 | 0.32 | 0.93 | 0.00 |

^{a,b} Different letters among breeds differ significantly ($P<0.05$).

¹ B=Black, G=Grey-Brown, R=Red-Brown, W=White and Y=Yellow-Brown.

Table 3. Color of the breast meats from five lines of Korean native chicken

| Line ¹ | L* | | a* | | b* | |
|-------------------|---------------------|--------|--------|--------------------|---------------------|---------------------|
| | Male | Female | Male | Female | Male | Female |
| B | 59.25 ^{ab} | 60.66 | 8.21 | 7.47 ^a | 20.70 ^c | 21.23 ^{bc} |
| G | 60.04 ^a | 60.29 | 7.29 | 7.00 ^{ab} | 21.67 ^a | 22.73 ^a |
| R | 59.07 ^{ab} | 60.55 | 8.26 | 6.88 ^{ab} | 21.38 ^{ab} | 21.94 ^b |
| W | 59.39 ^{ab} | 60.88 | 7.51 | 6.97 ^{ab} | 19.92 ^d | 20.80 ^c |
| Y | 58.62 ^b | 60.87 | 7.95 | 6.63 ^b | 20.83 ^{bc} | 21.57 ^b |
| SEM | 0.342 | 0.308 | 0.269 | 0.199 | 0.158 | 0.197 |
| | Line | Gender | Line | Gender | Line | Gender |
| <i>P</i> -value | 0.6433 | <.0001 | 0.0464 | <.0001 | <.0001 | <.0001 |
| <i>F</i> -value | 0.63 | 49.65 | 2.43 | 38.21 | 30.25 | 44.27 |

^{a~c} Different letters among breeds differ significantly ($P<0.05$).

¹ B=Black, G=Grey-Brown, R=Red-Brown, W=White, and Y=Yellow-Brown.

from male chickens of line B and the breast meat from female chickens of line W showed the lowest b* values ($P<0.05$). An effect of chicken line on the L*, a* and b* values of thigh meat was observed (Table 4). The lowest L* and highest a* values of thigh meat were found in male and female chickens of line Y ($P<0.05$). Additionally, thigh meat from male chickens of line Y had the lowest b* value ($P<0.05$). The lowest b* value for thigh meat from female chickens

was observed in line B ($P<0.05$).

An effect of sex on the L*, a* and b* values of both breast and thigh meat from the five KNC lines was observed. The breast and thigh meats of male chickens had lower L* and b* and higher a* values than those of female chickens (data not shown, $P<0.05$). Previous studies reported the sex effect on the color of chicken meat. However, the sex effect on color of chicken meat in the present study was not exactly

Table 4. Color of the thigh meats from five lines of Korean native chicken

| Line ¹ | L* | | a* | | b* | |
|-------------------|---------------------|---------------------|----------------------|---------------------|--------------------|---------------------|
| | Male | Female | Male | Female | Male | Female |
| B | 46.08 ^{bc} | 50.38 ^{ab} | 13.98 ^{abc} | 13.41 ^{ab} | 19.29 ^b | 20.76 ^b |
| G | 48.90 ^a | 52.00 ^{ab} | 13.24 ^c | 12.98 ^b | 20.36 ^a | 21.61 ^a |
| R | 47.13 ^{ab} | 51.28 ^a | 13.45 ^{bc} | 13.03 ^b | 19.52 ^b | 21.17 ^{ab} |
| W | 45.90 ^{bc} | 50.45 ^{ab} | 14.21 ^{ab} | 13.62 ^{ab} | 19.24 ^b | 20.97 ^{ab} |
| Y | 44.50 ^c | 50.37 ^b | 14.50 ^a | 14.03 ^a | 19.06 ^b | 21.27 ^{ab} |
| SEM | 0.468 | 0.404 | 0.242 | 0.202 | 0.195 | 0.172 |
| | Line | Gender | Line | Gender | Line | Gender |
| <i>P</i> -value | <.0001 | <.0001 | <.0001 | 0.0006 | <.0001 | <.0001 |
| <i>F</i> -value | 13.42 | 261.77 | 10.38 | 11.84 | 7.33 | 207.84 |

^{a~c} Different letters among breeds differ significantly ($P<0.05$).

¹ B=Black, G=Grey-Brown, R=Red-Brown, W=White and Y=Yellow-Brown.

coincide with that of previous studies. Schneider et al. (2012) reported a lower L^* and b^* and higher a^* values in breast meat from male chickens than in breast meat from female chickens in broiler at deboning. However, they also found no sex effect on a^* values of breast meat from broiler after 24 h postmortem. Souza et al. (2011) revealed that breast meat from male chickens had higher a^* and lower b^* values than did breast meat from female chickens of three broiler strains. López et al. (2011) reported a lower b^* value in breast meat from male chickens than in breast meat from female chickens in two broiler strains. Previous study reported that the difference of meat color between male and female chicken originated from different ultimate pH of meat related to the difference of glycogen content in muscle or regulatory patterns of post-mortem metabolism (Schneider et al., 2012). However, there was no significant difference in ultimate pH and WHC between meats from male and female chicken in the present study. Therefore, the sex effect on the color of chicken meat in the present study may be related to the difference of myoglobin content between meats from male and female chicken. Anadón (2002) reported that muscle of male chicken contained higher myoglobin, which was muscle pigment, than that of female chicken.

3. Correlation between the Measurement Traits of Chicken Meat

From the data pooled from each breast and thigh meat sample, the correlation between the measurement traits was analyzed (Table 5). The initial pH measured at 15 min post-mortem was negatively correlated (-0.11) with the a^* value of breast meat and positively correlated (0.09) with the b^* value of thigh meat. The ultimate pH of the meat was consistently correlated with the L^* value (negative) and a^* value (positive) of both breast and thigh meat. Low ultimate pH of meat is considered to result in pale meat color (Le Bihan-Duval et al. 2001). As explained above, low ultimate pH is strongly related to poor WHC in meat (Huff-Lonergan and Lonergan, 2005). However, a correlation (0.22) between the ultimate pH and WHC was only observed in thigh meat in the present study. It was previously reported that WHC was negatively correlated with L^* and b^* values and positively correlated with a^* value in chicken meat (Qiao et al., 2001).

Table 5. Correlation coefficient between the measurement traits of meats from five lines of Korean native chicken

| Line ¹ | Breast meat | | | Thigh meat | | |
|-------------------|---------------------|------------------------|---------|---------------------|------------------------|---------|
| | pH _{15min} | pH _{ultimate} | WHC | pH _{15min} | pH _{ultimate} | WHC |
| WHC | ns ¹ | ns | | ns | 0.22 | |
| L^* | ns | -0.24 | ns | ns | -0.12 | -0.12 |
| a^* | -0.11 | 0.18 | ns | ns | 0.13 | 0.21 |
| b^* | ns | ns | -0.12 | 0.09 | ns | ns |

¹ no significance ($P>0.05$).

In the present study, the WHC was negatively correlated with b^* value of breast meat, whereas no correlations were observed between WHC and the L^* and a^* values of breast meat. In addition, the L^* and a^* values of thigh meat showed negative and positive correlations with WHC, respectively. However, the b^* value of thigh meat was not correlated with WHC.

Le Bihan-Duval et al. (2008) reported significant heritability of pH, WHC and meat color in chicken and suggested that pH, WHC and color of chicken meat can be improved by selection. In the present study, the significant differences in initial pH at 15 min post-slaughter of the breast and thigh meat, and ultimate pH of the thigh meat were found in lines B, G, R, W and Y of KNC, although no significant difference in WHC was observed. Additionally, meat color was significantly different among the five KNC lines. Based on the results, we concluded that the line W in male chickens and the line G and W in female chickens may be good candidates for the selection to develop breed for high-quality meat because these lines showed property of high initial pH or/and ultimate pH. However, further investigation of the heritability of pH, WHC and meat color in all the five KNC lines is warranted.

적 요

재래닭고기는 broiler 닭고기와 비교하여 단백질 함량은 높고, 지방 함량은 낮을 뿐만 아니라, 독특한 풍미 및 조직감으로 인해 고품질 식육자원으로 여겨지고 있다. 게다가 최근 고유 유전자원 확보 및 유전자원의 다양성이 요구되고 있고, 고품질 식육에 대한 소비자의 요구가 증가함에 따라

재래닭을 이용한 새로운 고급 육질형 종계의 생산이 시도되고 있다. 따라서 본 연구는 깃털색에 따라 흑색, 회갈색, 적갈색, 백색 및 황갈색으로 분류되어 있는 한국 재래닭 5계통 닭고기의 품질 특성을 측정 비교하여 고급 육질형 종계 생산을 위한 선발 및 개량 육종 계획에 필요한 정보를 제공하고자 수행되었다. 본 연구 수행을 위해 한국 재래닭 5계통의 총 595 F₁ 개체를 20주령에 도계하였으며, 닭고기 품질 특성으로써 가슴육과 다리육의 도계 후 15분 pH, 최종 pH, 보수력 및 육색을 측정하여 5계통 간 차이를 비교하였고, 성별에 따른 차이 또한 비교하였다. 본 연구의 결과, 가슴육과 다리육의 도계 후 15분 pH는 계통 및 성별 간에 유의적인 차이가 있는 것으로 나타났다($P<0.05$). 하지만 가슴육과 다리육의 최종 pH 및 보수력 측정 결과, 다리육에서 계통 간에 최종 pH가 차이가 있었던 것을 제외하고는 계통 및 성별 간에 유의적인 차이가 없는 것으로 나타났다($P>0.05$). 육색 비교에서는 가슴육 L* 값을 제외하고, 가슴육과 다리육의 육색이 계통 및 성별 간에 유의적인 차이가 있는 것으로 확인되었다($P<0.05$). 측정된 품질 특성 간 상관관계 측정 결과, 최종 pH가 L* 값과 부의 상관관계를, 그리고 a* 값과 정의 상관관계를 가슴육과 다리육 모두에서 일괄되게 보임이 확인되었다($P<0.05$). 하지만 다른 품질 특성 간 상관관계는 가슴육과 다리육에서 일치하지 않았다. 따라서 본 연구의 결과, 계통 간 도계 후 15분 pH 및 최종 pH 차이를 고려하였을 때 고급 육질형 종계 생산을 위한 후보 계통으로서 수컷에서는 백색 계통이 암컷에서는 회갈색 및 백색 계통이 이용될 수 있을 것으로 사료된다. 하지만 추후 연구를 통해 재래닭고기의 도계 후 15분 pH, 최종 pH, 보수력 및 육색의 유전력 분석이 필요할 것으로 생각된다.

ACKNOWLEDGMENT

This research was supported by Golden Seed Project, Ministry of Agriculture, Food and Rural Affairs (MAFRA), Ministry of Oceans and Fisheries (MOF), Rural Development Administration (RDA), and Korea Forest Services (KFS). It was also partially supported by Research Institute of Eco-friendly Livestock Science, Institute of Green Bio Science and Technology, Seoul National University.

REFERENCES

Anadón HLS 2002 Biological, nutritional, and processing fac-

- tors affecting breast meat quality of broilers. Ph. D. Dissertation, Virginia Polytechnic Institute and State University, VA, USA.
- Allem CD, Russel SM, Fletcher DL 1997 The relationship of broiler breast meat color and pH to shelf-life and odor development. *Poultry Sci* 76:1042-1046.
- Barbut S, Sosnicki AA, Lonergan SM, Knapp T, Ciobanu DC, Gatcliffe LJ, Huff-Lonergan E, Wilson EW 2008 Progress in reducing the pale, soft and exudative (PSE) problem in pork and poultry meat. *Meat Sci* 79:46-63.
- Debut M, Berri C, Baéza E, Sellier N, Arnould C, Guémené D, Jehl N, Boutten B, Jégo Y, Beaumont C, Le Bihan-Duval E 2003 Variation of chicken technological meat quality in relation to genotype and preslaughter stress conditions. *Poultry Sci* 82:1829-1838.
- Dransfield E, Sosnicki AA 1999 Relationship between muscle growth and poultry meat quality. *Poultry Sci* 78:743-746.
- Fanatico AC, Pillai PB, Emmert JL, Owens CM 2007 Meat quality of slow- and fast-growing chicken genotypes fed low nutrient or standard diets and raised indoors or with outdoor access. *Poultry Sci* 86:2245-2255.
- Garcia RG, Freitas LWde, Schwingel AW, Farias RM, Caldara FR, Gabriel AMA, Graciano JD, Komiyama CM, Almeida Paz ICL 2010 Incidence and physical properties of PSE chicken meat in a commercial processing plant. *Braz J Poult Sci* 12:233-237.
- Huff-Lonergan E, Lonergan SM 2005 Mechanisms of water holding capacity in meat: The role of postmortem biochemical and structural changes. *Meat Sci* 71:194-204.
- Jaturasitha S, Srikanchai T, Kreuzer M, Wicke M 2008 Differences in carcass and meat characteristics between chicken indigenous to northern Thailand (Black-Boned and Thai native) and imported extensive breeds (Bresse and Rhode Island Red). *Poultry Sci* 87:160-169.
- Jayasena DD, Jung S, Kim HJ, Yong HI, Lee JH, Kim JG, Jo C 2013 Comparison of quality traits of meat from Korean native chickens and broilers used in two different traditional Korean cuisines. *Asian Australas J Anim Sci* 26:1038-1046.
- Jung S, Bae YS, Kim HJ, Jayasena DD, Lee JH, Park HB, Heo KN, Jo C 2013 Carnosine, anserine, creatine, and inosine 5'-monophosphate contents in breast and thigh meats

- from 5 lines of Korean native chicken. *Poultry Sci* 92: 3275-3282.
- Jung Y, Jeon HJ, Jung S, Choe JH, Lee JH, Heo KN, Kang BS, Jo C 2011 Comparison of quality traits of thigh meat from Korean native chickens and broilers. *Korean J Food Sci An* 31:684-692.
- Le Bihan-Duval E, Berri C, Baeza E, Millet N, Beaumont C 2001 Estimation of the genetic parameters of meat characteristics and of their genetic correlations with growth and body composition in an experimental broiler line. *Poultry Sci* 80:839-843.
- Le Bihan-Duval E, Debut M, Berri CM, Sellier N, Santé-Lhoutellier V, Jégo Y, Beaumont C 2008 Chicken meat quality: Genetic variability and relationship with growth and muscle characteristics. *BMG Genetics* 9:53-58.
- López KP, Schilling MW, Corzo A 2011 Broiler genetic strain and sex effects on meat characteristics. *Poultry Sci* 90: 1105-1111.
- Mancini RA, Hunt MC 2005 Current research in meat color. *Meat Sci* 71:100-121.
- Musa HH, Chen GH, Cheng HH, Shuiep ES, Bao WB 2006 Breed and sex effect on meat quality of chicken. *IJPS* 5: 566-568.
- Owens CM, Hirschler EM, McKee SR, Martinez-Dawson R, Sams AR 2000 The characterization and incidence of pale, soft, exudative turkey meat in a commercial plant. *Poultry Sci* 79:553-558.
- Qiao M, Fletcher DL, Smith DP, Northcutt JK 2001 The effect of broiler breast meat color on pH, moisture, water-holding capacity, and emulsification capacity. *Poultry Sci* 80:676-680.
- Schneider BL, Renema RA, Betti M, Carney VL, Zuidhof MJ 2012 Effect of holding temperature, shackling, sex, and age on broiler breast meat quality. *Poultry Sci* 91:468-477.
- Souza XR, Faria PB, Bressan MC 2011 Proximate composition and meat quality of broilers reared under different production systems. *Braz J Poult Sci* 13:15-20.
- Swatland HJ 2008 How pH causes paleness or darkness in chicken breast meat. *Meat Sci* 80:396-400.
- Zanetti E, De Marchi M, Dalvit C, Cassandro M 2010 Genetic characterization of local Italian breeds of chickens undergoing *in situ* conservation. *Poultry Sci* 89:420-427.

Received Jan. 30, 2015, Revised Feb. 17, 2015, Accepted Mar. 20, 2015