

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/271934462>

Potential of thermal image analysis for screening salt stress-tolerant soybean (*Glycine max*)

Article in *Plant Genetic Resources* · July 2014

DOI: 10.1017/S1479262114000422

CITATION

1

READS

85

4 authors, including:



Gyoung Ju Nah

Seoul National University

52 PUBLICATIONS 885 CITATIONS

[SEE PROFILE](#)



Do-Soon Kim

Seoul National University

79 PUBLICATIONS 776 CITATIONS

[SEE PROFILE](#)

Potential of thermal image analysis for screening salt stress-tolerant soybean (*Glycine max*)

Jin-Won Kim, Tae-Young Lee, Gyoungju Nah and Do-Soon Kim*

Department of Plant Science, Research Institute for Agriculture and Life Sciences, College of Agriculture and Life Sciences, Seoul National University, Seoul, Republic of Korea

Abstract

Non-destructive high-throughput phenotyping based on phenomics is an emerging technology for assessing the genetic diversity of various traits and screening in breeding programmes. In this study, non-destructive measurements of leaf temperature and chlorophyll fluorescence were conducted to investigate the physiological responses of soybean (*Glycine max*) to salt stress so as to set up a non-destructive screening method. Two-week-old seedlings of soybean in the V2 stage were treated with 0, 12.5, 25, 50 and 100 mM NaCl to induce salt stress. Three parameters, photosynthesis rate, stomatal conductance and chlorophyll fluorescence, decreased significantly, while soybean leaf temperature increased by exhibiting a positive correlation with NaCl concentration ($P < 0.001$). Soybean leaf temperature increased significantly at 50 mM NaCl when compared with the untreated control, although no visual symptom was observed. We selected leaf temperature as a major physiological parameter of salt stress as its measurement is much easier, faster and cheaper than that of other physiological parameters. Therefore, leaf temperature can be used for evaluating the responses to salt stress in soybean as a non-destructive and phenomic parameter. The results of this study suggest that non-destructive parameters such as chlorophyll fluorescence and leaf temperature are useful tools for assessing the genetic diversity of soybean with regard to salt stress tolerance and to screen salt stress-tolerant soybean for breeding.

Keywords: chlorophyll fluorescence; high-throughput screening; leaf temperature; phenomics; soybean

Introduction

Gene transformation technology and knowledge of genomics have allowed generating new genotypes more effectively in a short period of time. However, compared with our increasing capacity to generate new genotypes, our technologies for phenotyping and screening are still far behind to meet the needs of modern breeding due to the unavailability of high-throughput phenotyping methods under field conditions and high costs of field screening. The recent introduction of non-destructive phenotyping methods based on plant phenomics reveals technical possibilities of meeting the needs (Rascher

et al., 2011). Among the techniques introduced, thermal image and chlorophyll fluorescence analyses may be suitable for high-throughput screening (HTS) considering their implementation time and cost-effectiveness. Chen *et al.* (2005) used thermal image analysis to study drought stress responses in corn and soybean. The results of previous studies suggest that thermal image analysis can be used for screening water stress tolerance in genetic resources and breeding water stress-tolerant varieties.

Soybean (*Glycine max*) is known to be susceptible to salt stress (Katerji *et al.*, 2000). Screening salt tolerance in soybean can be done using non-destructive methods that measure phenomic responses of soybean in response to salt stress. Among the non-destructive methods based on phenomic responses, thermal image analysis may be the most effective one as thermal response is directly related to water status in plant. Therefore, this study demonstrated

*Corresponding author. E-mail: dosoonkim@snu.ac.kr

the feasibility of phenomic evaluation of salt stress responses in soybean by establishing a non-destructive method based on plant thermal images for screening salt tolerance.

Materials and methods

Seedlings of soybean (*G. max* cv. Daewon) at the second trifoliate (V2) stage were transplanted into 250 ml plastic pots. Treatment with 0, 12.5, 25, 50 and 100 mM NaCl was carried out to induce salt stress. The experiment consisted of a randomized block with four replicates. Photosynthesis rate, stomatal conductance (Li-COR 6400, Li-COR, Lincoln, Nebraska, USA) and chlorophyll contents (SPAD-502, Minolta, Tokyo, Japan) were measured in every leaflet in the first trifoliate stage and chlorophyll fluorescence (Handy PEA, Hansatech Instrument, King's Lynn, Norfolk, UK) was measured after 1 h of dark adaptation as ordinary physiological non-destructive parameters together with visual growth at 0, 1 and 2 d after NaCl treatment (DAT). Plant leaf temperature (T420, FLIR systems Inc., Täby, Sweden) was also measured at 0, 1 and 2 DAT. Thermal images were analysed using FLIR Tools 3.1 (FLIR systems Inc., USA) and MATLAB 8.1 (The MathWorks Inc., Natick, Massachusetts, USA) to determine soybean leaf temperature. All statistical analyses were carried out using SAS 9.3.

Results and discussion

Among the five physiological parameters, photosynthesis rate, stomatal conductance and chlorophyll fluorescence (F_v/F_m) were significantly decreased with an increase in NaCl concentration (Table S1, available online). A significant decrease was found at 50 and 100 mM NaCl concentrations from even 1 DAT, compared with the control, and it became much more significant at 2 DAT. No significant changes in chlorophyll contents and visual damage (data not shown) were observed until 2 DAT. Soybean leaf temperature increased significantly at 50 and 100 mM NaCl concentrations at 2 DAT (Fig. 1).

Chlorophyll fluorescence is widely used for estimating the quantum yield in photosynthesis *in vivo* (Genty *et al.*, 1989). Chlorophyll fluorescence values were significantly correlated with photosynthesis ($P < 0.001$; Table 1). Consequently, chlorophyll fluorescence can help determine the photosynthetic performance of soybean under salt stress.

Leaf temperature was significantly correlated with other parameters such as photosynthesis rate and stomatal conductance ($P < 0.001$; Table 1). Leaf temperature is closely related to photosynthetic performance including




Fig. 1. Thermal images (A) and leaf temperature (B) of soybean at 2 d after NaCl treatment. The vertical line indicates the standard error of four replications.

stomatal movement (Jossier *et al.*, 2010) and useful for estimating stomatal conductance (Jones, 1999) and transpiration rate (Katul *et al.*, 2000). Together, our findings and previous reports suggest leaf temperature to be an indicator of the whole plant status or physiological responses under salt stress.

Physiological parameters can be used for determining responses to salt stress within 2 DAT in soybean (Table S1, available online). Leaf temperature is a parameter that is easier and faster to measure than the other parameters, suggesting that it can be used for screening salt stress tolerance in soybean. Sirault *et al.* (2009) screened salt-tolerant wheat and barley using thermal image analysis. Furbank and Tester (2011) emphasized that thermal imaging can be applied to HTS to save time and cost and thus to improve phenotyping and crop breeding.

Compared with genotyping technology (e.g. next-generation sequencing), phenotyping technology is still far behind in terms of its speed and capacity (White *et al.*, 2012) and thus does not meet the technical demands of phenotyping-based screening. Fortunately, many efforts have been made using phenomic screening methods, the so-called next-generation phenotyping (Cobb *et al.*, 2013). Jones *et al.* (2009) quantified the response of rice and grapevine to water deficit using thermal infrared image analyses at the field level. Balota *et al.* (2008) demonstrated canopy temperature depression (CTD)

Table 1. Summary of correlation analysis results for non-destructive parameters of soybean treated with NaCl^a

	LT	CC	PR	SC	CF
LT	1	0.529*	0.850***	0.828***	0.689**
CC		1	0.660**	0.692**	0.517*
PR			1	0.931***	0.753**
SC				1	0.691**
CF					1

LT, leaf temperature; CC, chlorophyll contents; PR, photosynthesis rate; SC, stomatal conductance; and CF, chlorophyll fluorescence.

Values were significantly different: * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$.

^a The parameters were measured at 2 d after NaCl treatment.

during day and night and the relationship between CTD and other physiological and morphological traits, resulting in the confirmation of drought tolerance in wheat lines at the field level. To acquire more accurate information of water use efficiency, McAusland *et al.* (2013) combined thermal and chlorophyll fluorescence image information. Recently, field-based phenotyping has been implemented using a semi-automated technology (Comar *et al.*, 2012). Physiological and phenomic responses of soybean to salt stress and close correlations between thermal images and other parameters suggest that thermal image analysis can be used for assessing salt tolerance in soybean genetic resources and for screening salt-tolerant soybean. Although further studies are required to be carried out, thermal responses of soybean to salt stress can contribute to HTS in soybean breeding programmes.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1479262114000422>

Acknowledgements

This work was carried out with the support of “Cooperative Research Program for Agriculture Science & Technology Development (Project title: Development of High Throughput Screening (HTS) Technology based on Crop Phenomics, Project No. PJ00907801), Rural Development Administration, Republic of Korea.

References

Balota M, Payne WA, Evett SR and Peters TR (2008) Morphological and physiological traits associated with canopy temperature depression in three closely related wheat lines. *Crop Science* 48: 1897–1910.

- Chen D, Huang J and Jackson TJ (2005) Vegetation water content estimation for corn and soybeans using spectral indices derived from MODIS near- and short-wave infrared bands. *Remote Sensing of Environment* 98: 225–236.
- Cobb JN, DeClerck G, Greenberg A, Clark R and McCouch S (2013) Next-generation phenotyping: requirements and strategies for enhancing our understanding of genotype–phenotype relationships and its relevance to crop improvement. *Theoretical and Applied Genetics* 126: 867–887.
- Comar A, Burger P, de Solan B, Baret F, Daumard F and Hanocq J-F (2012) A semi-automatic system for high throughput phenotyping wheat cultivars in-field conditions: description and first results. *Functional Plant Biology* 39: 914–924.
- Furbank RT and Tester M (2011) Phenomics – technologies to relieve the phenotyping bottleneck. *Trends in Plant Science* 16: 635–644.
- Genty B, Briantais J-M and Baker NR (1989) The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica et Biophysica Acta (BBA) - General Subjects* 990: 87–92.
- Jones H (1999) Use of thermography for quantitative studies of spatial and temporal variation of stomatal conductance over leaf surfaces. *Plant, Cell & Environment* 22: 1043–1055.
- Jones HG, Serraj R, Loveys BR, Xiong L, Wheaton A and Price AH (2009) Thermal infrared imaging of crop canopies for the remote diagnosis and quantification of plant responses to water stress in the field. *Functional Plant Biology* 36: 978–989.
- Jossier M, Kroniewicz L, Dalmas F, Le Thiec D, Ephritikhine G, Thomine S, Barbier-Brygoo H, Vavasseur A, Filleur S and Leonhardt N (2010) The Arabidopsis vacuolar anion transporter, AtCLC, is involved in the regulation of stomatal movements and contributes to salt tolerance. *The Plant Journal* 64: 563–576.
- Katerji N, van Hoorn JW, Hamdy A and Mastroianni M (2000) Salt tolerance classification of crops according to soil salinity and to water stress day index. *Agricultural Water Management* 43: 99–109.
- Katul GG, Ellsworth DS and Lai CT (2000) Modelling assimilation and intercellular CO₂ from measured conductance: a synthesis of approaches. *Plant, Cell & Environment* 23: 1313–1328.
- McAusland L, Davey P, Kanwal N, Baker N and Lawson T (2013) A novel system for spatial and temporal imaging of intrinsic plant water use efficiency. *Journal of Experimental Botany* 64: 4993–5007.
- Rascher U, Blossfeld S, Fiorani F, Jahnke S, Jansen M, Kuhn AJ, Matsubara S, Martin LL, Merchant A and Metzner R (2011) Non-invasive approaches for phenotyping of enhanced performance traits in bean. *Functional Plant Biology* 38: 968–983.
- Sirault XR, James RA and Furbank RT (2009) A new screening method for osmotic component of salinity tolerance in cereals using infrared thermography. *Functional Plant Biology* 36: 970–977.
- White JW, Andrade-Sanchez P, Gore MA, Bronson KF, Coffelt TA, Conley MM, Feldmann KA, French AN, Heun JT, Hunsaker DJ, Jenks MA, Kimball BA, Roth RL, Strand RJ, Thorp KR, Wall GW and Wang G (2012) Field-based phenomics for plant genetics research. *Field Crops Research* 133: 101–112.